

Effects of pion potential and nuclear symmetry energy on the π^-/π^+ ratio in heavy-ion collisions at beam energies around the pion production threshold

Wen-Mei Guo^{1,2,3}, Gao-Chan Yong^{1,4,5,*}, Hang Liu⁶, and Wei Zuo^{1,4,5}

¹*Institute of Modern Physics,
Chinese Academy of Sciences,
Lanzhou 730000, China*

²*School of Physical Science and Technology,
Lanzhou University,
Lanzhou 730000, China*

³*University of Chinese Academy of Sciences,
Beijing 100049, China*

⁴*State Key Laboratory of Theoretical Physics,
Institute of Theoretical Physics,
Chinese Academy of Sciences,
Beijing, 100190*

⁵*Kavli Institute for Theoretical Physics,
Chinese Academy of Sciences,
Beijing 100190, China*

⁶*Texas Advanced Computing Center
(TACC) University of Texas at Austin,
Austin, Texas 78758, USA*

(Dated: May 12, 2015)

Abstract

Within the framework of the isospin-dependent Boltzmann-Uehling-Uhlenbeck (IBUU) transport model, we studied the effects of the pion potential and the symmetry energy on the pion production in the central $^{197}\text{Au} + ^{197}\text{Au}$ collisions around the pion production threshold. It is found that the pion potential has opposite effects on the value of π^-/π^+ ratio at low and high pion energies. The effect of the pion potential on the total π^-/π^+ ratio becomes larger in heavy-ion collisions at beam energies below the pion production threshold. And at beam energies below the pion production threshold, with the pion potential, the effect of the symmetry energy on the π^-/π^+ ratio becomes smaller compared with that above the pion production threshold.

PACS numbers: 25.70.-z, 21.65.Ef

*Electronic address: yonggaochan@impcas.ac.cn

I. INTRODUCTION

The density dependence of nuclear symmetry energy is crucial to understand the structure of exotic nuclei, dynamics of heavy-ion collisions, and many important issues in nuclear astrophysics such as neutron star cooling and supernova explosive [1–4]. The ratio π^-/π^+ was first proposed by *Li* as a sensitive observable for the high density behavior of the symmetry energy [5]. After that, a lot of studies on π^-/π^+ ratio and the symmetry energy were carried out [6–14]. However, recently different transport models, using the π^-/π^+ ratio to interpret the FOPI pion data [15], give contradict conclusions for the stiffness of the symmetry energy [16]. Such situation stimulates more detailed studies on pion production, such as different considerations of the medium effects on the pion production, including energy conservation on pion production [9, 17–19] and pion potential in medium [20]. The effects of pion potential [20–25] on pion production are in fact rarely studied in heavy-ion collision around the pion production threshold. It is thus necessary to investigate how the pion potential affects pion production since some experimental studies on pion production are doing at facilities that offer fast radioactive beams, such as the National Superconducting Cyclotron Laboratory (NSCL) and the Facility for Rare Isotope Beams (FRIB) in the USA, or the Radioactive Isotope Beam Facility (RIBF) in Japan.

II. THE IBUU MODEL AND THE PION POTENTIAL

In this study, we adopt the semi-classical transport model IBUU, in which an isospin- and momentum-dependent mean-field single nucleon potential is used [26], i.e.,

$$\begin{aligned}
 U(\rho, \delta, \mathbf{p}, \tau) = & A_u(x) \frac{\rho_{\tau'}}{\rho_0} + A_l(x) \frac{\rho_{\tau}}{\rho_0} \\
 & + B \left(\frac{\rho}{\rho_0} \right)^{\sigma} (1 - x\delta^2) - 8x\tau \frac{B}{\sigma + 1} \frac{\rho^{\sigma-1}}{\rho_0^{\sigma}} \delta \rho_{\tau'} \\
 & + \frac{2C_{\tau, \tau}}{\rho_0} \int d^3 \mathbf{p}' \frac{f_{\tau}(\mathbf{r}, \mathbf{p}')}{1 + (\mathbf{p} - \mathbf{p}')^2 / \Lambda^2} \\
 & + \frac{2C_{\tau, \tau'}}{\rho_0} \int d^3 \mathbf{p}' \frac{f_{\tau'}(\mathbf{r}, \mathbf{p}')}{1 + (\mathbf{p} - \mathbf{p}')^2 / \Lambda^2},
 \end{aligned} \tag{1}$$

where $\tau = 1/2(-1/2)$ for neutrons(protons), $\delta = (\rho_n - \rho_p)/(\rho_n + \rho_p)$ is the isospin asymmetry, and ρ_n, ρ_p denote neutron and proton densities, respectively. The parameter values $A_u(x), A_l(x), B, C_{\tau, \tau}, C_{\tau, \tau'}, \sigma$, and Λ can be found in Ref. [27]. $f_{\tau}(\mathbf{r}, \mathbf{p})$ is the phase-space




FIG. 1: (Color online) Density- and momentum-dependent symmetry potential as a function of nucleonic kinetic energy with parameter $x = 0, -1$.

distribution function at coordinate $\mathbf{r}$ and momentum $\mathbf{p}$. Different x parameters can be used to mimic different forms of the symmetry energy predicted by various many-body theories without changing any property of the symmetric nuclear matter and the symmetry energy at normal density. In fact, what is particularly interesting and important for nuclear reactions induced by neutron-rich nuclei is the isovector (symmetry) potential. The present used symmetry potential $U_{sym} = (U_n - U_p)/2\delta$, as shown in Figure 1, fits the Lane potential data quite well [28].

Pion-nucleus optical potential has been used to explain the existence of pionic atoms. A pion potential had been constructed by Toki et al., which can successfully describe the deeply bound states of pionic atoms [23]. And Jun Hong et al. constructed an isospin dependent pion potential [20] which writing as $U_{\pi\pm} = \pm 8S_{int0}\rho_T \frac{\rho_T^{\gamma-1}}{\rho_0^\gamma}$, where ρ_T is the isospin density $(\rho_p - \rho_n)/2$. This potential agrees well with Toki's pion optical potential [23], but it only represents the so called s-wave contribution to the π -nucleus potential. Shown in




FIG. 2: (Color online) Density- and momentum-dependent pion potential. The colored lines labelled by different momenta (MD) are Buss's potential [25]. The black solid line denotes momentum-independent (MID) potential given by Jun Hong et al. [20]. The left window shows negatively charged pion potential while the right window shows positively charged pion potential.




FIG. 3: (Color online) Isoscalar (Left) and isovector (Right) potentials of pion (only $\delta = 0.2$ case is shown) used in the IBUU model.

figure 2, the black solid line, which is labelled by MID is the pion potential constructed by Jun Hong et al. Here we choose the parameter $\gamma = 2$ and $S_{int0} = 20$ MeV [20]. It is seen that it is strongly density dependent, repulsive for the π^- and attractive for the π^+ .

Figure 2 also shows the other pion potential labelled by colored lines MD, including so-

called s-wave and p-wave contributions, which divides into three parts according to different regions of pionic momentum. In the low energy regime of $p_\pi < 80$ MeV, the results by Oset et al. are used [21, 22, 24]. When $p_\pi > 140$ MeV, we use the form of pion potential based on the Δ -hole model [25]. O. Buss also constructed an appropriate potential when $80 \text{ MeV} \leq p_\pi \leq 140 \text{ MeV}$ by an interpolating spline to gain a continuous derivative and therefore a continuous velocity of the pion [25]. From figure 2, we can clearly see that this potential is density- and momentum-dependent, is repulsive at low pionic momenta but becomes attractive at higher pionic momenta. It is noted from figure 2 that the “MID” pion potential is just the isovector potential whereas the “MD” pion potential includes both isoscalar and isovector potentials. The latter is clearly shown in figure 3. Because the isoscalar potential is overall positive and the isovector potential is positive for π^- while it is negative for π^+ , and also the “MD” isovector potential is overall not stronger than that of “MID” pion (isovector) potential, one sees the two pion potentials are quite different. In this article, based on the IBUU model, we use the second form of the pion potential to study effects of pion potential and nuclear symmetry energy in heavy-ion collisions at beam energies around the pion production threshold.

III. RESULTS AND DISCUSSIONS




FIG. 4: Left: Momentum distribution of π meson in the compression stage in the central $^{197}\text{Au} + ^{197}\text{Au}$ at $E_{beam} = 400$ MeV/nucleon. Right: Same as the Left, but for $E_{beam} = 200$ MeV/nucleon.

Since we use the density- and momentum-dependent pion potential, it is necessary to show the momentum distribution of π meson in heavy-ion collisions. Figure 4 shows the momentum distribution of π meson in the central $^{197}\text{Au} + ^{197}\text{Au}$ at $E_{beam} = 400$ and 200 MeV/nucleon, respectively. Because at the final stage of a reaction (the local baryon density of a π meson is zero) π suffers no pion potential, we in figure 4 just show the momentum distribution of π mesons in the compression stage. From the left panel of figure 4, it is shown that for the $^{197}\text{Au} + ^{197}\text{Au}$ at $E_{beam} = 400$ MeV/nucleon, π mesons are mainly located around $p = 58$ MeV/c and $p = 215$ MeV/c. While for $E_{beam} = 200$ MeV/nucleon case, π mesons are mainly located around $p = 51$ MeV/c. In the compressed stage, pion in the nuclear matter can be absorbed by surrounding matter at certain momentum of pion meson, thus one can see a sunken shape of pion distribution in momentum space. This sunken shape is mainly decided by the kinematics of pion as well as its absorption mechanism in nuclear matter while it is less affected by the mean-field potential of pion. Different momentum distributions of the π mesons at different incident beam energies should cause different effects of pion potential on the charged pion ratio.




FIG. 5: (Color online) Left: Effect of pion potential on charged pion ratio π^-/π^+ in the central $^{197}\text{Au} + ^{197}\text{Au}$ at $E_{beam} = 400$ MeV/nucleon with the softer symmetry energy ($x = 0$). Right: Same as the Left, but for the stiffer symmetry energy ($x = -1$).

Figure 5 shows the effect of pion potential on charged pion ratio π^-/π^+ . It is seen that whether for the softer symmetry energy or the stiffer symmetry energy, around the Coulomb peak [7], the value of π^-/π^+ ratio with pion potential is higher than that without pion potential. But for energetic pion mesons, a reversed case occurs. The reason is that for π^- with low energy, π^- meson suffers a repulsive pion potential (as shown in figure 2). π^- mesons are less absorbed thus more π^- production. Due to the Coulomb action, π^+ mesons are less affected by the pion potential. Therefore we see a high value of π^-/π^+ ratio around the Coulomb peak. For the energetic π^- mesons, they suffers an attractive pion potential thus more π^- mesons are absorbed by surrounding nuclear matter. Then a low value of π^-/π^+ ratio is seen.




FIG. 6: (Color online) Effect of the symmetry energy on the π^-/π^+ ratio in the central $^{197}\text{Au} + ^{197}\text{Au}$ at $E_{beam} = 400$ MeV/nucleon without (Left) and with (Right) the pion potential.

To answer whether the effect of the symmetry energy on the π^-/π^+ ratio is affected by the pion potential, we plot figure 6. It is seen that the effect of the symmetry energy on the π^-/π^+ ratio is not affected much by the pion potential at incident beam energy of 400 MeV/nucleon. This is more clearly shown in the figure 7, time evolution of the π^-/π^+ ratio with and without pion potential for softer ($x=0$) and stiffer ($x=-1$) symmetry energies. From the figure 7, we see that total π^-/π^+ ratio is clearly affected by the symmetry energy.




FIG. 7: (Color online) Time evolution of the π^-/π^+ ratio with and without pion potential for the softer ($x=0$) and the stiffer ($x=-1$) symmetry energies.

The pion potential has less effect on the π^-/π^+ ratio. As pointed out in the figure 4, at incident beam energy of 400 MeV/nucleon, π mesons are mainly located around $p=58$ MeV/c and $p=215$ MeV/c. Therefore some pion mesons suffer repulsive pion potential, but others suffer attractive pion potential (shown in the figure 2). This counterbalance makes pion potential has less effect on the π^-/π^+ ratio at $E_{beam}=400$ MeV/nucleon, thus the effect of the symmetry energy on π^-/π^+ ratio is kept.

Since such related experiments are doing at NSCL/MSU ($E_{beam} < 150$ MeV/nucleon) and RIKEN/Japan ($E_{beam} = 200 \sim 350$ MeV/nucleon), it is necessary to show effects of pion potential and nuclear symmetry energy on the π^-/π^+ ratio in heavy-ion collisions at beam energies below the pion production threshold. Shown in figure 8 is the effect of pion potential on the π^-/π^+ ratio at the incident beam energy of 200 MeV/nucleon with the softer ($x=0$) and the stiffer ($x=-1$) symmetry energies. Again, around the Coulomb peak and in high pionic energy region the pion potential affects the π^-/π^+ ratio much. And the effect of pion potential on the π^-/π^+ ratio at $E_{beam}=200$ MeV/nucleon seems larger than that at $E_{beam}=400$ MeV/nucleon. This is understandable since at $E_{beam}=200$ MeV/nucleon, pion mesons are only located around $p=51$ MeV/c where only a repulsive pion potential dominates. To




FIG. 8: (Color online) Same as Figure 5, but for $E_{beam}=200$ MeV/nucleon.

see how the large effect of pion potential on the π^-/π^+ ratio at such beam energy affect the probing of the symmetry energy by the π^-/π^+ ratio, we plot figure 9, i.e., effect of the symmetry energy on the π^-/π^+ ratio without and with pion potential. It is seen that with pion potential, effect of the symmetry energy on the π^-/π^+ ratio almost disappears. This is also clearly seen in the figure 10, time evolution of the π^-/π^+ ratio with and without pion potential for the softer ($x=0$) and the stiffer ($x=-1$) symmetry energies. As demonstrated in Ref. [20], the large asymmetry $(\rho_n - \rho_p)/\rho$ of nuclear matter causes a large pion potential. A larger pion potential for pion mesons with lower energy causes π^- mesons are less absorbed by surrounding matter thus more π^- mesons are produced finally. However, on the other hand the large asymmetry of the nuclear matter also causes large repulsion to neutrons by the symmetry potential, thus less neutron-neutron collisions in matter make less π^- mesons production. Therefore, the pion potential cancels out the effect of symmetry energy on the π^-/π^+ ratio in heavy-ion collisions at beam energies below the pion production threshold.




FIG. 9: (Color online) Same as Figure 6, but for $E_{beam} = 200$ MeV/nucleon.

IV. CONCLUSIONS

In conclusions, we studied the effects of the pion potential and the symmetry energy on the π^-/π^+ ratio in the central $^{197}\text{Au} + ^{197}\text{Au}$ collisions around the pion production threshold. Because the symmetry potential in nuclear matter generally repels neutrons out of the dense matter, which causes small number of the neutron-neutron collisions. Small number of neutron-neutron collisions causes small number of π^- to produce. However, the pion potential generally repels π^- out of the dense matter thus π^- 's are less absorbed. In heavy-ion collisions, the pion potential causes large number of π^- 's whereas the symmetry potential causes small number of π^- 's. The effects of the symmetry energy on the π^-/π^+ ratio thus cancelled out by the pion potential in some degree.

Acknowledgments

The author G.C. Yong thanks Prof. Eulogio Oset and Dr. Jun Hong for communications. The authors acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing HPC resources that have contributed to the research results reported within this paper. This work is supported by the National Natural Science




FIG. 10: (Color online) Same as Figure 7, but for $E_{beam} = 200$ MeV/nucleon.

Foundation of China (11375239, 11435014, 11175219). The 973 Program of China under No. 2013CB834405, the Knowledge Innovation Project (KJCX2-EW-N01) of Chinese Academy of Sciences.

-
- [1] B.A. Li, L.W. Chen, and C.M. Ko, Phys. Rep. **464**, 113 (2008).
 - [2] V. Baran, M. Colonna, V. Greco, M. Di Toro, Phys. Rep. **410**, 335 (2005).
 - [3] J.M. Lattimer, M. Prakash, Science **304**, 536 (2004).
 - [4] A.W. Steiner, M. Prakash, J.M. Lattimer, et al., Phys. Rep. **411**, 325 (2005).
 - [5] B.A. Li, Phys. Rev. Lett. **88**, 192701 (2002).
 - [6] B.A. Li, G.C. Yong, W. Zuo, Phys. Rev. C **71**, 014608 (2005).
 - [7] G.C. Yong, B.A. Li, L.W. Chen, W. Zuo, Phys. Rev. C **73**, 034603 (2006).
 - [8] Q.F. Li, Z. Li, S. Soff, M. Bleicher, H. Stöcker, J. Phys. G **32**, 151 (2006).
 - [9] G. Ferini et al., Phys. Rev. Lett. **97**, 202301 (2006); G. Ferini et al., Nucl. Phys. A **762**, 147 (2005).
 - [10] Y. Gao, L. Zhang, H.F. Zhang, X.M. Chen, G.C. Yong, Phys. Rev. C **83**, 047602 (2011).
 - [11] Y. Gao, G.C. Yong, Y.J. Wang, Q.F. Li, W. Zuo, Phys. Rev. C **88**, 057601 (2013).

- [12] Z.G. Xiao, B.A. Li, L.W. Chen, G.C. Yong, M. Zhang, Phys. Rev. Lett. **102**, 062502 (2009).
- [13] Z.Q. Feng, G.M. Jin, Phys. Lett. B **683**, 140 (2010).
- [14] W.J. Xie, J. Su, L. Zhu, F.S. Zhang, Phys. Lett. B **718**, 1510 (2013).
- [15] W. Reisdorf et al., Nucl. Phys. A **781**, 459 (2007).
- [16] W.M. Guo, G.C. Yong, Y.J. Wang, Q.F. Li, H.F. Zhang, W. Zuo, Phys. Lett. B **738**, 397 (2014).
- [17] J. Xu, L.W. Chen, C.M. Ko, B.A. Li, Y.G. Ma, Phys. Rev. C **87**, 067601 (2013).
- [18] C.M. Ko, Y. Oh, and J. Xu, Int. J. Mod. Phys. E **19**, 1763 (2010).
- [19] M.D. Cozma, arXiv:1409.3110 (2014).
- [20] J. Hong and P. Danielewicz, Phys. Rev. C **90**, 024605 (2014).
- [21] M. Ericson and T.E. O. Ericson, Ann. of Phys. **36**, 323 (1966).
- [22] C. García-Recio, E. Oset and L.L. Salcedo, Phys. Rev. C **37**, 194 (1988).
- [23] H. Toki et al., Nucl. Phys. A **501**, 653 (1989).
- [24] J. Nieves, E. Oset, C. Garcia-Recio, Nucl. Phys. A **554**, 554 (1993).
- [25] O. Buss, diploma thesis, <https://gibuu.hepforge.org/trac/wiki/Paper#Diplomatheses> (2004).
- [26] C.B. Das et al., Phys. Rev. C **67**, 034611 (2003).
- [27] B.A. Li, C.B. Das, S. Das Gupta, and C. Gale, Nucl. Phys. A **735**, 563 (2004); Phys. Rev. C **69**, 064602 (2004).
- [28] L.W. Chen, C.M. Ko, B.A. Li and G.C. Yong, Frontiers of Physics in China **2**, 327 (2007).